

DESCRIPTION

PROCESS FOR PRODUCING 5-HYDROXY-4-THIOMETHYLPYRAZOLE
COMPOUND

5

TECHNICAL FIELD

The present invention relates to a process for producing a 5-hydroxy-4-thiomethylpyrazole compound, which is useful as an intermediate in the production of pharmaceutical and agricultural chemicals.

10

BACKGROUND ART

The 5-hydroxy-4-thiomethylpyrazole compound obtained by the present invention is useful as an intermediate in the production of pharmaceutical and agricultural chemicals.

15

As for the process for producing a 4-thiomethylpyrazole compound, a method of reacting a 4-chloromethylpyrazole compound and a sulfur compound is known (see, Patent Document 1).

20

In order to obtain a 4-chloromethylpyrazole compound used as the raw material of this reaction, for example, after once synthesizing a pyrazole compound having a methyl group at the 4-position by a ring-closing reaction of corresponding β -ketoesters with hydrazines, the methyl group must be converted into a chloromethyl group by chlorination. Furthermore, for obtaining a 4-thiomethylpyrazole compound, this chloromethyl group must be reacted with an organic sulfur compound (see, Patent Document 1). However, this method is not satisfied as the industrial process for producing a 4-thiomethylpyrazole compound, because the reaction comprises multiple stages and the yield is not high due to difficulty in the selective monohalogenation of the methyl group at the 4-position of pyrazole.

25

30

35

As for the process for producing the 4-chloromethylpyrazole compound as the raw material in the

above-described method, a method of directly
chloromethylating a pyrazole compound unsubstituted at
the 4-position is also known (see, Non-Patent Document
1). However, this method is problematic and hardly
5 employable as the industrial production method, because
the reaction is associated with by-production of a
bis(chloromethyl) ether which is a carcinogenic
substance.

Furthermore, a reaction of directly thiomethylating
10 a pyrazole compound is not known.

(Patent Document 1) International Patent
Publication WO2004/013106

(Non-Patent Document 1) Journal of chemical
Society, pp. 1205-1208 (1955)

DISCLOSURE OF THE INVENTION

A method capable of solving those drawbacks in
conventional techniques and simply and easily producing a
5-hydroxy-4-thiomethylpyrazole compound in a good yield
20 has been demanded.

Under these circumstances, the present inventors
have made intensive studies on the process for producing
a 5-hydroxy-4-thiomethylpyrazole compound, as a result,
it has been found that surprisingly, when a 5-
25 hydroxypyrazole compound represented by formula (1) shown
later is reacted with a sulfur compound represented by
formula (2) shown later in the presence of a base and
formaldehyde, a 5-hydroxy-4-thiomethylpyrazole compound
represented by formula (3) shown later is produced. The
30 present invention has been accomplished based on this
finding.

According to the process of the present invention
having such a constitution, a 5-hydroxy-4-
thiomethylpyrazole compound represented by formula (3)
35 can be produced from a 5-hydroxypyrazole compound
represented by formula (1) in a good yield through a
single step by a simple and easy operation method under


mild conditions without using a special reaction apparatus or an expensive catalyst or transition metal. Moreover, in the process of the present invention, a harmful waste derived from a catalyst or transition metal is substantially not produced and therefore, this process is environment-friendly and has high industrial utility value.

In addition, according to the embodiment of using water as the solvent in the process of the present invention, a more environment-friendly process having a higher industrial utility value is provided.
(Embodiment of the Invention)

The present invention includes, for example, the following embodiments [1] to [6].

[1] A process for producing a 5-hydroxy-4-thiomethylpyrazole compound, comprising: reacting a pyrazole represented by formula (1):

(Chemical Formula 1)



(1)

(wherein R_1 represents a hydrogen atom, an alkyl group, an aromatic hydrocarbon group capable of having a substituent, or a heterocyclic group capable of having a substituent, and R_2 represents an electron-withdrawing group) with a sulfur compound represented by formula (2):


(Chemical Formula 2)



(wherein X represents a hydrogen atom or a metal, R_3 represents an alkyl group, an aromatic hydrocarbon group capable of having a substituent, or a heterocyclic group capable of having a substituent, and n represents 0 or 2) in the presence of a base and formaldehyde to produce a 5-hydroxy-4-thiomethylpyrazole compound represented by

formula (3):

(Chemical Formula 3)



(wherein R_1 , R_2 , R_3 and n have the same meanings as above).

[2] The process for producing a 5-hydroxy-4-thiomethylpyrazole compound as described in [1], wherein n is 0.

[3] The process for producing a 5-hydroxy-4-thiomethylpyrazole compound as described in [1], wherein n is 2.

[4] The process for producing a 5-hydroxy-4-thiomethylpyrazole compound as described in any one of [1] to [3], wherein the electron-withdrawing group represented by R_2 is a trifluoromethyl group.

[5] The process for producing a 5-hydroxy-4-thiomethylpyrazole compound as described in any one of [1] to [3], wherein the electron-withdrawing group represented by R_2 is a cyano group.

[6] The process for producing a 5-hydroxy-4-thiomethylpyrazole compound as described in any one of [1] to [3], wherein the electron-withdrawing group represented by R_2 is an alkoxycarboxyl group, a carboxyl group or a metal salt thereof.

BEST MODE FOR CARRYING OUT THE INVENTION

The present invention is described in detail below. In the following, unless otherwise indicated, the "parts" and "%" denoting a quantitative ratio are on the mass basis.

(Process for producing 5-Hydroxy-4-Thiomethylpyrazole Compound)

The present invention relates to a process for producing a 5-hydroxy-4-thiomethylpyrazole compound represented by formula (3), comprising reacting a 5-hydroxypyrazole compound represented by formula (1) with
5 a sulfur compound represented by formula (2) in the presence of a base and formaldehyde.
(Isomer)

The 5-hydroxypyrazole compound represented by formula (1) used as the raw material in the present
10 invention and the 5-hydroxy-4-thiomethyl pyrazole compound represented by formula (3) as the product may be present in the form of a keto-enol tautomer. In the present invention, the raw material compound and the product are represented by a structure based on the enol
15 form as in formulae (1) and (3), but in practice, the compositional ratio of the keto-enol tautomer sometimes varies depending on, for example, the solvent for dissolving the compound and even in such a case, the raw material or product in the process of the present
20 invention includes both isomers of keto form/enol form.
(5-Hydroxypyrazole Compound)

The 5-hydroxypyrazole compound represented by formula (1) used as the raw material in the present invention is described below.

25 The method for obtaining the 5-hydroxypyrazole compound represented by formula (1) is not particularly limited. That is, any one method in the following examples may be used or other methods may also be used.
(Examples of Synthesis Method)

30 With respect to the synthesis method of the 5-hydroxypyrazole compound, for example, Hiroshi Yamanaka et al., Hetero-Kan Kagobutsu no Kagaku (Chemistry of Heterocyclic Compounds), Chap. 5, Kodansha Scientific (1988), and J.A. Joule and K. Mills, Handbook of
35 Heterocyclic Chemistry, 2nd ed., Chap. 4.3.2.3, Pergamon (2000) may be referred to.

For example, a method of reacting a corresponding β -

ketoester compound with a hydrazine, more specifically, a method where 1-methyl-5-hydroxy-3-trifluoromethylpyrazole can be synthesized in a yield of 49% by refluxing ethyl 4,4,4-trifluoroacetoacetate and methylhydrazine with a water solvent under heating for 2 hours, is reported in Journal of Heterocyclic Chemistry, Vol. 27, page 243 (1990).

Similarly, a method of reacting an oxaloacetic diester with a hydrazine to obtain a 3-(alkoxycarbonyl)-5-hydroxypyrazole compound, and a method of converting the alkoxycarbonyl group of the obtained compound into a cyano group are described in detail in Kokai (Japanese Unexamined Patent Publication) No. 10-287654.

Also, a method for obtaining a 3-cyano-5-hydroxypyrazole compound by a reaction of an α -cyanosuccinic acid with a diazonium salt is described in Kokoku (Japanese Examined Patent Publication) No. 51-33556.
(Substituent R_1)

As for formula (1) representing a 5-hydroxypyrazole compound which is the raw material compound in the process of the present invention, examples of the substituent represented by R_1 in formula (1) include the followings:

(1) a hydrogen atom;

(2) a linear or branched C1-C6 alkyl group having a carbon number of 1 to 6 (hereinafter, as for the carbon number, for example, when the carbon number is from 1 to 6, this is simply referred to as "C1-C6"), such as methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-butyl group, n-pentyl group and n-hexyl group;

(3) a monocyclic or condensed-ring aromatic hydrocarbon group with the number of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as phenyl group and naphthyl group; the aromatic hydrocarbon group may have one or more substituent such

as those in the following (3.1) to (3.21):

(3.1) a halogen atom such as bromo, chloro, fluoro and iodo,

5 (3.2) a linear or branched C1-C6 alkyl group such as methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-butyl group, n-pentyl group and n-hexyl group,

(3.3) a hydroxyl group,

10 (3.4) a linear or branched C1-C6 alkoxy group such as methoxy group, ethoxy group, n-propoxy group and isopropoxy group,

(3.5) a linear or branched C1-C6 hydroxyalkyl group such as hydroxymethyl group and 1-hydroxyethyl group,

15 (3.6) a linear or branched (C1-C6 alkoxy)-(C1-C6 alkyl) group such as methoxymethyl group, 1-methoxyethyl group and 1-ethoxyethyl group,

(3.7) a linear or branched C1-C6 haloalkyl group such as fluoromethyl group, difluoromethyl group and 20 trifluoromethyl group,

(3.8) a carboxyl group,

(3.9) a metal salt of a carboxyl group, as represented by an alkali metal salt such as sodium salt, potassium salt and lithium salt, and an alkaline earth 25 metal salt such as calcium salt, barium salt and magnesium salt,

(3.10) a linear or branched (C1-C6 alkoxy)carbonyl group such as methoxycarbonyl group and ethoxycarbonyl group,

30 (3.11) an arylcarbonyl group with the number of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as benzoyl group and naphthoyl group,

(3.12) a monocyclic or condensed-ring 35 heteroarylcarbonyl group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number

- of atoms constituting the ring being from 5 to 14,
preferably from 5 to 10, such as pyridylcarbonyl group,
thienylcarbonyl group and furylcarbonyl group,
- (3.13) a nitro group,
- 5 (3.14) an amino group,
- (3.15) a linear or branched mono- or di-(C1-C6
alkyl)amino group such as methylamino group,
dimethylamino group, ethylamino group and diethylamino
group,
- 10 (3.16) a linear or branched (C1-C6
alkyl)carbonylamino group such as acetylamino group,
propionylamino group and butyrylamino group,
- (3.17) a linear or branched hydroxycarbonyl (C1-C6
alkyl) group such as hydroxycarbonylmethyl group and 1-
15 hydroxycarbonylethyl group,
- (3.18) a linear or branched (C1-C6
alkoxy)carbonyl-(C1-C6 alkyl) group such as
methoxycarbonylmethyl group, 1-methoxycarbonylethyl group
and 1-ethoxycarbonylethyl group,
- 20 (3.19) a linear or branched aminocarbonyl-(C1-C6
alkyl) group such as aminocarbonylmethyl group and 1-
aminocarbonylethyl group,
- (3.20) a linear or branched (C1-C6
alkyl)aminocarbonyl-(C1-C6 alkyl) group such as
25 methylaminocarbonylmethyl group, 1-
methylaminocarbonylethyl group and 1-
ethylaminocarbonylethyl group, and
- (3.21) a cyano group;
- (4) a monocyclic or condensed-ring aromatic
30 heterocyclic group having from 1 to 4 heteroatoms which
are at least one member selected from a nitrogen atom, an
oxygen atom and a sulfur atom, with the number of atoms
constituting the ring being from 5 to 14, preferably from
5 to 10, as represented by, for example, a furyl group, a
35 benzofuryl group, a pyridyl group, a thienyl group, a
benzothienyl group, an oxazolyl group, a benzoxazolyl
group, a thiazolyl group, a benzothiazolyl group, an

isooxazolyl group, a thiadiazolyl group, a pyrazyl group, a pyrimidinyl group, an indolyl group, a quinolinyl group, a pyrazolyl group, an imidazolyl group, a benzimidazolyl group, a triazolyl group and a triazinyl group; the aromatic heterocyclic group may have one or more substituent such as those in the following (4.1) to (4.19):

(4.1) a linear or branched C1-C6 alkyl group such as methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-butyl group, n-pentyl group and n-hexyl group,

(4.2) a hydroxyl group,

(4.3) a linear or branched C1-C6 alkoxy group such as methoxy group, ethoxy group, n-propoxy group and isopropoxy group,

(4.4) a linear or branched C1-C6 hydroxyalkyl group such as hydroxymethyl group and hydroxyethyl group,

(4.5) a linear or branched (C1-C6 alkoxy)-(C1-C6 alkyl) group such as methoxymethyl group, methoxyethyl group and ethoxyethyl group,

(4.6) a linear or branched C1-C6 haloalkyl group such as fluoromethyl group, difluoromethyl group and trifluoromethyl group,

(4.7) a carboxyl group,

(4.8) a metal salt of a carboxyl group, as represented by an alkali metal salt such as sodium salt, potassium salt and lithium salt, and an alkaline earth metal salt such as calcium salt, barium salt and magnesium salt,

(4.9) a linear or branched (C1-C6 alkoxy)carbonyl group such as methoxycarbonyl group and ethoxycarbonyl group,

(4.10) a halogen atom such as bromo, chloro, fluoro and iodo,

(4.11) a nitro group,

(4.12) an amino group,

(4.13) a linear or branched mono- or di-(C1-C6

alkyl)amino group such as methylamino group, dimethylamino group, ethylamino group and diethylamino group,

5 (4.14) a linear or branched (C1-C6 alkyl)carbonylamino group such as acetylamino group, propionylamino group and butyrylamino group,

(4.15) a cyano group,

(4.16) a formyl group,

10 (4.17) a linear or branched (C1-C6 alkyl)carbonyl group such as methylcarbonyl group and ethylcarbonyl group,

(4.18) an arylcarbonyl group with the number of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as benzoyl group and naphthoyl group, and

15 (4.19) a monocyclic or condensed-ring heteroarylcarbonyl group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, such as pyridylcarbonyl group, thienylcarbonyl group and furylcarbonyl group; and

(5) a monocyclic or condensed-ring non-aromatic heterocyclic group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, as represented by, for example, a hydrofuryl group, a pyranlyl group, a thioranyl group, a thianlyl group, a pyrrolidinyl group, an indolinyl group, a piperidinyl group, an imidazolidinyl group and a piperazinyl group; the non-aromatic heterocyclic group may have one or more substituent such as those in the following (5.1) to (5.19):

35 (5.1) a linear or branched C1-C6 alkyl group such as methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-

butyl group, n-pentyl group and n-hexyl group,

(5.2) a hydroxyl group,

(5.3) a linear or branched C1-C6 alkoxy group
such as methoxy group, ethoxy group, n-propoxy group and
5 isopropoxy group,

(5.4) a linear or branched C1-C6 hydroxyalkyl
group such as hydroxymethyl group and hydroxyethyl group,

(5.5) a linear or branched (C1-C6 alkoxy)-(C1-C6
alkyl) group such as methoxymethyl group, methoxyethyl
10 group and ethoxyethyl group,

(5.6) a linear or branched C1-C6 haloalkyl group
such as fluoromethyl group, difluoromethyl group and
trifluoromethyl group,

(5.7) a carboxyl group,

15 (5.8) a metal salt of a carboxyl group, as
represented by an alkali metal salt such as sodium salt,
potassium salt and lithium salt, and an alkaline earth
metal salt such as calcium salt, barium salt and
magnesium salt,

20 (5.9) a linear or branched (C1-C6
alkoxy)carbonyl group such as methoxycarbonyl group and
ethoxycarbonyl group,

(5.10) a halogen atom such as bromo, chloro,
fluoro and iodo,

25 (5.11) a nitro group,

(5.12) an amino group,

(5.13) a linear or branched mono- or di-(C1-C6
alkyl)amino group such as methylamino group,
dimethylamino group, ethylamino group and diethylamino
30 group,

(5.14) a linear or branched (C1-C6
alkyl)carbonylamino group such as acetylamino group,
propionylamino group and butyrylamino group,

(5.15) a cyano group,

35 (5.16) a formyl group,

(5.17) a linear or branched (C1-C6 alkyl)carbonyl
group such as methylcarbonyl group and ethylcarbonyl

group,

(5.18) an arylcarbonyl group with the number of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as benzoyl group and naphthoyl group, and

(5.19) a monocyclic or condensed-ring heteroarylcarbonyl group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, such as pyridylcarbonyl group, thienylcarbonyl group and furylcarbonyl group.
(Preferred Substituent R₁)

Among those described above, in view of easy availability, the followings are suitably usable as the substituent R₁.

A C1-C6 alkyl group as represented by methyl, and an aromatic hydrocarbon group with the number of atoms constituting the ring being from 6 to 14, as represented by phenyl are preferred, and a methyl group and a phenyl group are more preferred.
(Electron-Withdrawing Group R₂)

The electron-withdrawing group represented by R₂ in formula (1) means an atomic group capable of withdrawing an electron from the other party by the induction effect, an aromatic hydrocarbon group having such an atomic group, or an aromatic heterocyclic group having such an atomic group. Specific examples of the electron-withdrawing group R₂ include the followings:

(1) a linear or branched C1-C6 haloalkyl group such as difluoromethyl group and trifluoromethyl group; a carboxyl group, or a metal salt of a carboxyl group, as represented by an alkali metal salt such as sodium salt, potassium salt and lithium salt, and an alkaline earth metal salt such as calcium salt, barium salt and magnesium salt;

(2) a linear or branched (C1-C6 alkoxy)carbonyl

group such as methoxycarbonyl group and ethoxycarbonyl group;

(3) a halogen atom such as bromo, chloro, fluoro and iodo; a nitro group; a formyl group;

5 (4) a linear or branched (C1-C6 alkyl)carbonyl group such as methylcarbonyl group (acetyl group) and ethylcarbonyl group;

(5) an arylcarbonyl group with the number of atoms constituting the ring being from 6 to 14, preferably from
10 6 to 10, such as benzoyl group and naphthoyl group;

(6) a monocyclic or condensed-ring heteroarylcarbonyl group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number
15 of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, such as pyridylcarbonyl group, thienylcarbonyl group and furylcarbonyl group;

(7) aminocarbonyl group, a linear or branched mono- or di-(C1-C6 alkyl)aminocarbonyl group such as
20 methylaminocarbonyl group and dimethylaminocarbonyl group; a cyano group;

(8) an aromatic hydrocarbon group containing, as the substituent, one or more atomic group capable of withdrawing an electron from the other party by the
25 induction effect, with the number of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as chlorophenyl group (e.g., 2,4-dichlorophenyl), carboxyphenyl group and nitrophenyl group; and

(9) a monocyclic or condensed-ring aromatic
30 heterocyclic group containing, as the substituent, one or more atomic group capable of withdrawing an electron from the other party by the induction effect, and having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom,
35 with the number of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, such as chlorofuryl group, chlorobenzofuryl group, chloropyridyl group,

chlorothienyl group, carboxylbenzothienyl group,
nitrooxazolyl group, cyanobenzooxazolyl group,
chlorothiazolyl group, chlorobenzothiazolyl group, acetyl
isooxazolyl group, benzoylthiadiazolyl group,
5 chloropyrazyl group, chloropyrimidinyl group,
chloroindolyl group, chloroquinolinyl group,
aminocarbonylpyrazolyl group,
methylaminocarbonylimidazolyl group,
trifluoromethylbenzoimidazolyl group,
10 methoxycarbonyltriazolyl group, and chlorotriazinyl
group.

(Preferred Electron-Withdrawing Group R_2)

Among those described above, in view of stability
under the reaction condition, that is, in the presence of
15 a base, and eventual stability in the yield, the
followings are suitably usable as the electron-
withdrawing group R_2 .

A C1-C6 haloalkyl group as represented by
trifluoromethyl; an aromatic hydrocarbon group
20 containing, as the substituent, one or more atomic group
capable of withdrawing an electron from the other party
by the induction effect, with the number of atoms
constituting the ring being from 6 to 14, as represented
by 2,4-dichlorophenyl; and a cyano group are preferred,
25 and a trifluoromethyl group, 2,4-dichlorophenyl and a
cyano group are more preferred.

(Preferred 5-Hydroxypyrazole Compound)

Accordingly, specific examples of the 5-
hydroxypyrazole compound represented by formula (1)
30 include 5-hydroxy-3-trifluoromethylpyrazole, 3-
ethoxycarbonyl-5-hydroxypyrazole, 3-acetyl-5-
hydroxypyrazole, 3-benzoyl-5-hydroxypyrazole, 5-hydroxy-
3-(3-pyridylcarbonyl)pyrazole, 3-cyano-5-hydroxypyrazole,
5-hydroxy-1-methyl-3-trifluoromethylpyrazole, 3-
35 ethoxycarbonyl-5-hydroxy-1-methylpyrazole, 3-chloro-5-
hydroxy-1-methylpyrazole, 5-hydroxy-1-methyl-3-
nitropyrazole, 5-hydroxy-1-methyl-3-(2-

thienylcarbonyl)pyrazole, 5-hydroxy-1-methyl-3-(3-pyridylcarbonyl)pyrazole, 3-dimethylaminocarbonyl-5-hydroxy-1-methylpyrazole, 5-hydroxy-1-methyl-3-(4-trifluoromethylphenyl)pyrazole, 3-(4-ethoxycarbonylphenyl)-5-hydroxy-1-methylpyrazole, 3-(2,4-dichlorophenyl)-5-hydroxy-1-methylpyrazole, 3-(3,5-dinitrophenyl)-5-hydroxy-1-methylpyrazole, 3-(4-dimethylaminocarbonyl)-5-hydroxy-1-methylphenylpyrazole, 5-hydroxy-1-n-propyl-3-trifluoromethylpyrazole, 3-cyano-1-n-hexyl-5-hydroxypyrazole, 1-tert-butyl-5-hydroxy-3-trifluoromethylpyrazole, 1-tert-butyl-3-(4-carboxyphenyl)-5-hydroxypyrazole, 3-(4-acetylphenyl)-1-tert-butyl-5-hydroxypyrazole, 1-tert-butyl-3-(4-cyanophenyl)-5-hydroxypyrazole, 5-hydroxy-1-phenyl-3-trifluoromethylpyrazole, 3-cyano-5-hydroxy-1-phenylpyrazole, 1-(4-chlorophenyl)-3-ethoxycarbonyl-5-hydroxypyrazole, 3-ethoxycarbonyl-5-hydroxy-1-(2-methylphenyl)pyrazole, 3-ethoxycarbonyl-5-hydroxy-1-(2-methoxymethylphenyl)pyrazole, 1-(4-acetylphenyl)-3-ethoxycarbonyl-5-hydroxypyrazole, 3-ethoxycarbonyl-5-hydroxy-1-(3-nitrophenyl)pyrazole, 5-hydroxy-1-(2-methoxyphenyl)-3-trifluoromethylpyrazole, 5-hydroxy-3-trifluoromethyl-1-(4-trifluoromethylphenyl)pyrazole, 1-(4-ethoxycarbonylphenyl)-5-hydroxy-3-trifluoromethylpyrazole, 1-(4-dimethylaminophenyl)-5-hydroxy-3-trifluoromethylpyrazole, 1-(4-acetylaminophenyl)-5-hydroxy-3-trifluoromethylpyrazole, 1-(4-methoxycarbonylmethylphenyl)-5-hydroxy-3-trifluoromethylpyrazole, 1-(4-dimethylaminocarbonylmethylphenyl)-5-hydroxy-3-trifluoromethylpyrazole, 1-(4-cyanophenyl)-5-hydroxy-3-trifluoromethylpyrazole, 1-(2-naphthyl)-5-hydroxy-3-trifluoromethylpyrazole, 1-(2-benzothiazolyl)-5-hydroxy-3-trifluoromethylpyrazole, 5-hydroxy-1-(2-pyridyl)-3-trifluoromethylpyrazole and 5-hydroxy-1-(2-pyrimidyl)-3-trifluoromethylpyrazole.

(Sulfur Compound)

The sulfur compound represented by formula (2) is described below.

Examples of the substituent represented by R_3 in formula (2) include the followings:

- 5 (1) a linear or branched C1-C6 alkyl group such as methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-butyl group, n-pentyl group and n-hexyl group; a monocyclic or condensed-ring aromatic hydrocarbon group with the number
10 of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as phenyl group and naphthyl group (the aromatic hydrocarbon group may have one or more substituent, for example, a linear or branched C1-C6 alkyl group such as methyl group, ethyl
15 group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-butyl group, n-pentyl group and n-hexyl group,
- (2) a hydroxyl group,
- (3) a linear or branched C1-C6 alkoxy group such as
20 methoxy group, ethoxy group, n-propoxy group and isopropoxy group,
- (4) a linear or branched C1-C6 hydroxyalkyl group such as hydroxymethyl group and hydroxyethyl group,
- (5) a linear or branched (C1-C6 alkoxy)-(C1-C6
25 alkyl) group such as methoxymethyl group, methoxyethyl group and ethoxyethyl group,
- (6) a linear or branched C1-C6 haloalkyl group such as fluoromethyl group, difluoromethyl group and trifluoromethyl group, a carboxyl group, a metal salt of
30 a carboxyl group, as represented by an alkali metal salt such as sodium salt, potassium salt and lithium salt, and an alkaline earth metal salt such as calcium salt, barium salt and magnesium salt,
- (7) a linear or branched (C1-C6 alkoxy)carbonyl
35 group such as methoxycarbonyl group and ethoxycarbonyl group,
- (8) a halogen atom such as bromo, chloro, fluoro

and iodo, a nitro group, an amino group,

5 (9) a linear or branched mono- or di-(C1-C6 alkyl)amino group such as methylamino group, dimethylamino group, ethylamino group and diethylamino group,

(10) a linear or branched C1-C6 alkylcarbonylamino group such as acetylamino group, propionylamino group and butyrylamino group, a cyano group, a formyl group,

10 (11) a linear or branched (C1-C6 alkyl)carbonyl group such as methylcarbonyl group and ethylcarbonyl group,

(12) an arylcarbonyl group with the number of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as benzoyl group and naphthoyl group, and

15 (13) a monocyclic or condensed-ring heteroarylcarbonyl group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, such as pyridylcarbonyl group, thienylcarbonyl group and furylcarbonyl group); and

20 (14) an aromatic or non-aromatic heterocyclic group including a monocyclic or condensed-ring aromatic heterocyclic group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, as represented by, for example, a furyl group, a benzofuryl group, a pyridyl group, a thienyl group, a benzothienyl group, an oxazolyl group, a benzoxazolyl group, a thiazolyl group, a benzothiazolyl group, an isooxazolyl group, a thiadiazolyl group, a pyrazyl group, a pyrimidinyl group, an indolyl group, a quinolinyl group, a pyrazolyl group, an imidazolyl group, a benzimidazolyl group, a triazolyl group and a triazinyl group (the aromatic heterocyclic group may have one or more substituent, for example, a linear or branched C1-C6

25

30

35

alkyl group such as methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, sec-butyl group, tert-butyl group, n-pentyl group and n-hexyl group,

(15) a hydroxyl group,

5 (16) a linear or branched C1-C6 alkoxy group such as methoxy group, ethoxy group, n-propoxy group and isopropoxy group,

(17) a linear or branched C1-C6 hydroxyalkyl group such as hydroxymethyl group and hydroxyethyl group,

10 (18) a linear or branched (C1-C6 alkoxy)-(C1-C6 alkyl) group such as methoxymethyl group, methoxyethyl group and ethoxyethyl group,

(19) a linear or branched C1-C6 haloalkyl group such as fluoromethyl group, difluoromethyl group and
15 trifluoromethyl group,

(20) a carboxyl group, or a metal salt of a carboxyl group, as represented by an alkali metal salt such as sodium salt, potassium salt and lithium salt, and an alkaline earth metal salt such as calcium salt, barium
20 salt and magnesium salt,

(21) a linear or branched (C1-C6 alkoxy)carbonyl group such as methoxycarbonyl group and ethoxycarbonyl group,

(22) a halogen atom such as bromo, chloro, fluoro
25 and iodo, a nitro group, an amino group,

(23) a linear or branched mono- or di-(C1-C6 alkyl)amino group such as methylamino group, dimethylamino group, ethylamino group and diethylamino group,

30 (24) a linear or branched (C1-C6 alkyl)carbonylamino group such as acetylamino group, propionylamino group and butyrylamino group, a cyano group, a formyl group,

(25) a linear or branched (C1-C6 alkyl)carbonyl group such as methylcarbonyl group and ethylcarbonyl
35 group,

(26) an arylcarbonyl group with the number of atoms constituting the ring being from 6 to 14, preferably from

6 to 10, such as benzoyl group and naphthoyl group, and
(27) a monocyclic or condensed-ring
heteroarylcarbonyl group having from 1 to 4 heteroatoms
which are at least one member selected from a nitrogen
5 atom, an oxygen atom and a sulfur atom, with the number
of atoms constituting the ring being from 5 to 14,
preferably from 5 to 10, such as pyridylcarbonyl group,
thienylcarbonyl group and furylcarbonyl group), and a
monocyclic or condensed-ring non-aromatic heterocyclic
10 group having from 1 to 4 heteroatoms which are at least
one member selected from a nitrogen atom, an oxygen atom
and a sulfur atom, with the number of atoms constituting
the ring being from 5 to 14, preferably from 5 to 10, as
represented by, for example, a hydrofuryl group, a
15 pyranlyl group, a thioranlyl group, a thianlyl group, a
pyrrolidinyl group, an indolinyl group, a piperidinyl
group, an imidazolidinyl group and a piperazinyl group
(the non-aromatic heterocyclic group may have one or more
substituent, for example, a linear or branched C1-C6
20 alkyl group such as methyl group, ethyl group, n-propyl
group, isopropyl group, n-butyl group, sec-butyl group,
tert-butyl group, n-pentyl group and n-hexyl group, a
hydroxyl group,
(28) a linear or branched C1-C6 alkoxy group such as
25 methoxy group, ethoxy group, n-propoxy group and
isopropoxy group,
(29) a linear or branched C1-C6 hydroxyalkyl group
such as hydroxymethyl group and hydroxyethyl group,
() a linear or branched (C1-C6 alkoxy)-(C1-C6
30 alkyl) group such as methoxymethyl group, methoxyethyl
group and ethoxyethyl group,
(30) a linear or branched C1-C6 haloalkyl group such
as fluoromethyl group, difluoromethyl group and
trifluoromethyl group, a carboxyl group, a metal salt of
35 a carboxyl group, as represented by an alkali metal salt
such as sodium salt, potassium salt and lithium salt, and
an alkaline earth metal salt such as calcium salt, barium

salt and magnesium salt,

(31) a linear or branched (C1-C6 alkoxy)carbonyl group such as methoxycarbonyl group and ethoxycarbonyl group,

5 (32) a halogen atom such as bromo, chloro, fluoro and iodo, a nitro group, an amino group,

(33) a linear or branched mono- or di-(C1-C6 alkyl)amino group such as methylamino group, dimethylamino group, ethylamino group and diethylamino group,

10 (34) a linear or branched (C1-C6 alkyl)carbonylamino group such as acetylamino group, propionylamino group and butyrylamino group, a cyano group, a formyl group,

(35) a linear or branched (C1-C6 alkyl)carbonyl group such as methylcarbonyl group and ethylcarbonyl group,

15 (36) an arylcarbonyl group with the number of atoms constituting the ring being from 6 to 14, preferably from 6 to 10, such as benzoyl group and naphthoyl group, and

20 (37) a monocyclic or condensed-ring heteroarylcarbonyl group having from 1 to 4 heteroatoms which are at least one member selected from a nitrogen atom, an oxygen atom and a sulfur atom, with the number of atoms constituting the ring being from 5 to 14, preferably from 5 to 10, such as pyridylcarbonyl group, thienylcarbonyl group and furylcarbonyl group).

(Group X)

X in formula (2) represents, for example, a hydrogen atom; or a metal atom as represented by, for example, an alkali metal such as sodium, potassium and lithium, and an alkaline earth metal such as magnesium and calcium.

25 In the case where n in formula (2) is 0, the compound is a thiol or a salt thereof, and when n is 2, the compound is a sulfinic acid or a salt thereof.

35 In the sulfur compound represented by formula (2) for use in the reaction, X may be a hydrogen atom or a metal salt of an alkali metal such as sodium, potassium

and lithium, or an alkaline earth metal such as magnesium and calcium. Furthermore, in the case where the sulfur compound is a thiol (a compound when n in formula (2) is 0), a precursor capable of producing a thiol, as
5 represented by isothiuronium salt or the like which can be easily prepared by a reaction of the corresponding alkyl halide with thiourea, may be hydrolyzed in the same reaction vessel, so that a thiol working out to the raw material in the reaction can be formed in the system and
10 used.

(Specific Example of Sulfur Compound)

Accordingly, specific examples of the sulfur compound represented by formula (2) include sodium
thiomethoxide, sodium thioethoxide, 2-butanethiol,
15 thiophenol, 2-ethylthiophenol, 4-methoxythiophenol, 4-chlorothiophenol, 4-nitrothiophenol, 4-dimethylaminothiophenol, 4-cyanothiophenol, 4-acetylthiophenol, 2-mercaptopyridine, 2-mercaptobenzoxazole, 2-mercaptobenzothiazole, an
20 isothiuronium salt as represented by n-hexyl thiocarboxamidine hydrochloride, benzyl thiocarboxamidine hydrochloride and [5,5-dimethyl(4,5-dihydroisooxazol-3-yl)]thiocarboxamidine hydrochloride, sodium benzenesulfinate and sodium p-toluenesulfinate.

25 (Reaction Method)

The method of reacting a 5-hydroxypyrazole compound represented by formula (1) with a sulfur compound represented by formula (2) to produce a 5-hydroxy-4-thiomethylpyrazole compound represented by formula (3) is
30 described below.

(Formaldehyde)

The above-described reaction is performed in the presence of formaldehyde. The form of formaldehyde for use in this reaction is not particularly limited and
35 formaldehyde in any form may be used, but use of an aqueous formaldehyde solution at a concentration of 35 to 50% as represented by 35% formalin easily available as a

commercial product, or paraformaldehyde (a polymer of formaldehyde, which produces formaldehyde in the system by undergoing hydrolysis and therefore, is usable as a formaldehyde equivalent) is easy in view of operation and preferred.

(Amount Used)

The amount of formaldehyde used may be sufficient if it is an equivalent amount or more per equivalent of the raw material compound represented by formula (1), but the amount used is usually from 1.0 to 5.0 equivalents, preferably from 1.0 to 3.0 equivalents, per mol of the raw material compound represented by formula (1).

The amount used of the sulfur compound represented by formula (2) may be sufficient if it is an equivalent amount or more to the raw material compound of formula (1), but the amount used is usually from 1.0 to 2.0 equivalents, preferably from 1.0 to 1.2 equivalents, per mol of the raw material compound represented by formula (1).

(Base)

In the present invention, the reaction is performed in the presence of a base. Examples of the base for use in the reaction include the followings:

(1) an alkali metal hydride such as sodium hydride, potassium hydride and lithium hydride;

(2) an alkali metal such as sodium metal, potassium metal and lithium metal;

(3) an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide and lithium hydroxide;

(4) an alkaline earth metal hydroxide such as barium hydroxide, magnesium hydroxide and calcium hydroxide;

(5) an alkali metal carbonate such as sodium carbonate, potassium carbonate, sodium hydrogencarbonate and potassium hydrogencarbonate;

(6) an inorganic base as represented by, for example, an alkaline earth metal oxide such as barium

oxide, magnesium oxide and calcium oxide; and a metal alkoxide such as sodium methoxide, sodium ethoxide, potassium methoxide, potassium ethoxide and tert-butoxy potassium; and

- 5 (7) an organic base as represented by, for example, an alkali metal such as butyllithium.

Among these, an alkali metal hydroxide and a metal alkoxide are preferred in view of availability, and an alkali metal hydroxide, particularly, sodium hydroxide,
10 is more preferred, because the reaction can be performed with a water solvent and therefore, the load such as wastewater aftertreatment can be reduced.

(Amount of Base Used)

The amount of the base used may be any amount as
15 long as it is large enough to allow for satisfactory progress of the reaction, but the amount used is, for example, from 1.0 to 20 mol, preferably from 1.5 to 10 mol, more preferably from 1.5 to 3.0 mol, per mol of the 5-hydroxypyrazole compound (raw material compound)
20 represented by formula (1).

(Solvent)

The reaction of the present invention may be performed in the presence of a solvent, if desired.

The solvent usable for the reaction may be
25 sufficient if it does not inhibit the reaction, and examples thereof include water; alcohols such as methanol and ethanol; aromatic hydrocarbons such as toluene, xylene and chlorobenzene; halogenated aliphatic hydrocarbons such as dichloromethane and chloroform;
30 aprotic polar solvents such as dimethylformamide, dimethylacetamide, N-methylpyrrolidone, tetramethylurea, hexamethylphosphoric triamide (HMPA) and propylene carbonate; ether-based solvents such as diethyl ether, tetrahydrofuran and dioxane; and aliphatic hydrocarbons
35 such as pentane and n-hexane. These solvents may be used individually or as a mixed solvent at an arbitrary mixing ratio. In view of solubility and reactivity of the base,

the reaction is preferably performed by using water or alcohols, more preferably in water or methanol.

5 The amount of the solvent may be sufficient if it is large enough to allow for satisfactory stirring of the reaction system, but the amount thereof is usually from 0.05 to 10 L (liter), preferably from 0.5 to 2 L, per mol of the 5-hydroxypyrazole compound (raw material compound) represented by formula (1).

(Reaction Conditions)

10 The reaction temperature of this reaction may be, for example, from 0°C to the reflux temperature of the solvent used, but the reaction is preferably performed at 20 to 50°C, particularly, at room temperature with stirring, because this is simple and a good yield is
15 obtained.

The reaction time of this reaction is not particularly limited, but the reaction can be satisfactorily completed usually in the time from 1 to 10 hours.

20 According to this reaction, a 5-hydroxy-4-thiomethylpyrazole compound represented by formula (3) can be produced in a good yield by a simple and easy operation method under mild conditions. The obtained 5-hydroxy-4-thiomethylpyrazole compound represented by
25 formula (3) is a compound useful as an intermediate raw material for pharmaceutical and agricultural chemicals.

(EXAMPLES)

30 The process for producing the compound of the present invention is specifically described below by referring to Examples, but the present invention is not limited to these Examples.

(Reference Example 1)

35 Synthesis of 5-hydroxy-1-methyl-3-trifluoromethylpyrazole:

Ethyl 4,4,4-trifluoroacetoacetate (92.1 g (0.5 mol))

was dissolved in 60.1 g (1.0 mol) of acetic acid. The resulting solution was cooled to 10°C or less with stirring, and 65.8 g (0.5 mol) of an aqueous 35% methylhydrazine solution was added dropwise thereto over 1 hour. After the dropwise addition, the solution was stirred at room temperature for 1 hour and subsequently at 80°C for 5 hours. When the reaction was completed, the reaction solution was cooled to room temperature, and 150 mL (milliliter) of toluene, 600 mL of water and 48 g (1.2 mol) of sodium hydroxide were added thereto. After liquid separation, 154 g (1.5 mol) of 35% hydrochloric acid was added dropwise to the obtained aqueous layer, and the produced crystal was collected by filtration. The crystal was washed twice with 50 mL of water and dried by a hot air drier to obtain 71.8 g (yield: 86.5%) of the title compound as a pale yellow crystal.

LC-MS (EI): $m/z=166$ (M^+), melting point: 179-180°C.

(Reference Example 2)

Synthesis of 5-hydroxy-1-phenyl-3-trifluoromethylpyrazole:

Ethyl 4,4,4-trifluoroacetoacetate (18.4 g (0.1 mol)) was dissolved in 12.0 g (0.2 mol) of acetic acid. The resulting solution was cooled to 10°C or less with stirring, and 11.8 g (0.11 mol) of phenylhydrazine was added dropwise thereto over 0.5 hours. After the dropwise addition, the solution was stirred at room temperature for 1 hour and subsequently at 80°C for 5 hours. When the reaction was completed, the reaction solution was cooled to room temperature, and 100 mL of water was added thereto. The produced crystal was collected by filtration, washed twice with 50 mL of water and dried by a hot air drier to obtain 22.3 g (yield: 98.0%) of the title compound as a pale yellow crystal.

LC-MS(EI): $m/z=228$ (M^+), melting point: 190-192°C.

(Reference Example 3)

Synthesis of 5-hydroxy-3-trifluoromethylpyrazole hydrochloride:

5 Ethyl 4,4,4-trifluoroacetoacetate (18.4 g (0.1 mol))
was dissolved in 20 mL of ethanol. The resulting
solution was cooled to 10°C or less with stirring, and 6.0
g (0.12 mol) of hydrazine was added dropwise thereto over
0.5 hours. After the dropwise addition, the solution was
10 stirred at room temperature for 1 hour and subsequently
at the reflux temperature for 5 hours. When the reaction
was completed, the reaction solution was cooled to room
temperature, and 100 mL of water and 20.6 g (0.2 mol) of
35% hydrochloric acid were added thereto. The produced
crystal was collected by filtration, washed twice with 10
15 mL of water and dried by a hot air drier to obtain 12.8 g
(yield: 68.1%) of the title compound as a white crystal.

LC-MS(EI): m/z=152 (M⁺).

(Reference Example 4)

20 Synthesis of 3-ethoxycarbonyl-5-hydroxy-1-methylpyrazole:

Monosodium diethyl oxaloacetate (50.0 g (0.24 mol))
was suspended in 500 mL of ethanol, and 25 mL of acetic
acid was added thereto. Subsequently, 15 g (0.33 mol) of
97% methylhydrazine was added dropwise thereto over 0.5
25 hours with stirring. After the dropwise addition, the
solution was stirred at room temperature for 2 hours and
then at the reflux temperature for 5 hours. The
resulting solution was cooled, ethanol was removed by
distillation under reduced pressure, and 200 mL of ethyl
30 acetate and 100 mL of water were added to the residue.
After liquid separation, the aqueous layer was re-
extracted with 50 mL of ethyl acetate, and the combined
ethyl acetate layer was washed with 50 mL of water and
then with 50 mL of saturated brine. The obtained ethyl
35 acetate layer was dried over anhydrous sodium sulfate,
and the solvent was removed by distillation under reduced
pressure. After adding 100 mL of water to the

precipitated crystal, the crystal was collected by filtration, washed with 10 mL of water and dried by a hot air drier to obtain 29.2 g (yield: 71.8%) of the title compound as a pale yellow crystal.

5 LC-MS(EI): $m/z=170$ (M^+), 125 (base), melting point: 151°C.

(Reference Example 5)

Synthesis of 3-cyano-5-hydroxy-1-phenylpyrazole:

10 Water (120 mL) and 15 mL of 35% hydrochloric acid were added to 5.6 g (0.06 mol) of aniline, and the mixture was dissolved. Subsequently, 24 mL of water having dissolved therein 4.2 g (0.06 mol) of sodium
15 nitrite was added dropwise thereto with stirring under ice cooling to 0 to 5°C, and the resulting solution was stirred for 1 hour to prepare benzenediazonium chloride. An aqueous solution of this diazonium salt was added dropwise to a 120 mL pyridine solution containing 10.2 g
20 (0.06 mol) of diethyl α -cyanosuccinate with stirring under ice cooling. After the dropwise addition, the solution was stirred under ice cooling for 1 hour and then at room temperature for 1 hour. When the reaction was completed, 240 mL of an aqueous 2% sodium hydroxide solution was added, followed by stirring for 2 hours.
25 The obtained reaction solution was added dropwise to 240 mL of 35% hydrochloric acid under ice cooling, and the precipitated crystal was collected by filtration, washed with 10 mL of water and dried by a hot air drier to obtain 8.4 g of the crude crystal of the title compound
30 as a red-brown crystal. This crude crystal was recrystallized from diethyl ether-petroleum ether and dried by a hot air drier to obtain 5.7 g (yield: 51.3%) of the title compound as a pale yellow crystal.

35 LC-MS(EI): $m/z=185$ (M^+), 125 (base), melting point: 190°C.

(Reference Example 6)

Synthesis of 3-(2,4-dichlorophenyl)-5-hydroxy-1-methylpyrazole:

Potassium ethyl malonate (22.1 g (0.13 mol)) was
5 suspended in 200 mL of ethyl acetate, and 29.14 g (0.29
mol) of triethylamine and 13.7 g (0.14 mol) of magnesium
chloride were sequentially added thereto with stirring
under ice cooling. The resulting suspension liquid was
stirred at 40°C for 6 hours and again ice-cooled, and
10 20.9 g (0.1 mol) of 2,4-dichlorobenzoyl chloride was
added dropwise thereto over 1 hour while keeping the
reaction solution at 10°C or less. After the dropwise
addition, the stirring was continued at room temperature
for 12 hours. Subsequently, 200 mL of 5% hydrochloric
15 acid was added dropwise to the reaction solution, and the
organic layer was separated. The obtained organic layer
was washed twice with 50 mL of water and then with 30 mL
of saturated brine, and the solvent was removed by
distillation under reduced pressure. To the residue
20 obtained by the concentration, 100 mL of ethanol was
added. The resulting solution was cooled to 10°C or less
with stirring, and 13.1 g (0.1 mol) of an aqueous 35%
methylhydrazine solution was added dropwise thereto over
1 hour. After the dropwise addition, the solution was
25 stirred at room temperature for 1 hour and then at 80°C
for 3 hours. When the reaction was completed, the
reaction solution was cooled to room temperature, 300 mL
of water was added, and the produced crystal was
collected by filtration. The crystal was washed twice
30 with 50 mL of water and dried by a hot air drier to
obtain 12.3 g (yield: 50.2%) of the title compound as a
white crystal.

^1H -NMR value (300 MHz, CDCl_3): σ =7.53 (d; J =1.8 Hz,
1H), 7.2-7.4 (m, 2H), 5.68 (s, 1H), 3.54 (s, 3H) ppm.

35 LC-MS (EI): m/z =242 [$(M-1)^+$], melting point: 221-
223°C.

(Reference Example 7)

Synthesis of 3-(3,5-dinitrophenyl)-5-hydroxy-1-methylpyrazole:

5 Potassium ethyl malonate (11.0 g (0.07 mol)) was
suspended in 100 mL of ethyl acetate, and 14.5 g (0.19
mol) of triethylamine and 6.9 g (0.08 mol) of magnesium
chloride were sequentially added thereto with stirring
under ice cooling. The resulting suspension liquid was
10 stirred at 40°C for 6 hours and again ice-cooled, and
11.5 g (0.05 mol) of 3,5-dinitrobenzoyl chloride was
added dropwise thereto over 1 hour while keeping the
reaction solution at 10°C or less. After the dropwise
addition, the stirring was continued at room temperature
15 for 12 hours. Subsequently, 100 mL of 5% hydrochloric
acid was added dropwise to the reaction solution, and the
organic layer was separated. The obtained organic layer
was washed twice with 50 mL of water and then with 30 mL
of saturated brine, and the solvent was removed by
20 distillation under reduced pressure. To the residue
obtained by the concentration, 100 mL of ethanol was
added, and 2.4 g (0.05 mol) of 97% methylhydrazine was
then added dropwise at room temperature with stirring.
After the dropwise addition, the solution was stirred at
25 room temperature for 1 hour and subsequently at 80°C for 3
hours. When the reaction was completed, the reaction
solution was cooled to room temperature, 300 mL of water
was added, and the produced crystal was collected by
filtration. The crystal was washed twice with 50 mL of
30 water, and the obtained crystal was recrystallized from
50% aqueous ethanol and dried by a hot air drier to
obtain 4.2 g (yield: 30.2%) of the title compound as a
red crystal.

¹H-NMR value (300 MHz, MeOH-d₄): σ =8.90 (t; J=2.1 Hz,
35 1H), 8.85 (d; J=2.1 Hz, 2H), 4.60 (s, 1H), 3.73 (s, 3H),
3.3-3.4 (m, 1H) ppm.

LC-MS (EI): $m/z=264$ (M^+), melting point: 230-231°C.

(Reference Example 8)

Synthesis of 3-chloro-5,5-dimethyl-4,5-dihydroisooxazole:

5 Ethanol (500 mL) and 63.0 g (0.75 mol) of sodium
hydrogencarbonate were added and stirred at room
temperature. After blowing 84.2 g (1.50 mol) of
isobutene gas for 0.5 hours, the temperature was elevated
to 70°C, and 131.3 g (0.5 mol) of a 40% isopropyl ether
10 solution of dichloroformoxime was gradually added
dropwise to the reaction solution, followed by stirring
at the same temperature for 8 hours. The resulting
reaction solution was allowed to cool to 25°C or less and
after removing inorganic solids by filtration, the
15 residue was subjected to distillation under reduced
pressure of 62°C/1.1 kPa to obtain 32.3 g (yield: 51%) of
3-chloro-4,5-dihydroisooxazole as a colorless transparent
liquid.

20 ^1H -NMR value (300 MHz, MeOH-d_4): $\sigma=2.88$ (s, 2 H),
1.41 (s, 3H) ppm.

LC-MS (EI): $m/z=133$ (M^+), 118 (base), boiling point:
50°C/0.7 kPa.

(Reference Example 9)

25 Synthesis of [5,5-dimethyl(4,5-dihydroisooxazol-3-yl)]-
thiocarboxamidine hydrochloride:

35% Hydrochloric acid (4.17 g (0.04 mol)) was added
to an ethanol 100 mL (0.51/mol) solution containing 16.8
g (0.2 mol) of thiourea, and 26.7 g (0.2 mol) of 3-
30 chloro-5,5-dimethyl-4,5-dihydroisooxazole was added
dropwise thereto over 1 hour with stirring at room
temperature, followed by stirring at 30°C for 3 hours.
Subsequently, 100 mL of toluene was added to the reaction
solution and when the solvent was removed by distillation
under reduced pressure, 58.3 g of a white crystal was
35 precipitated. Thereafter, 300 mL of isopropyl alcohol

was added to the obtained crude crystal, and the mixture was heated until the crystal was dissolved, and then gradually cooled to obtain 38.5 g (yield: 92.0%) of the title compound as a white crystal.

5 ¹H-NMR value (300 MHz, MeOH-d₄): σ=4.84 (s, 3 H), 3.08 (s, 2H), 1.46 (s, 3H) ppm.

Melting point: 147°C.

10 Example 1: Synthesis of 5-hydroxy-1-methyl-4-methylthiomethyl-3-trifluoromethylpyrazole

5-Hydroxy-1-methyl-3-trifluoromethylpyrazole (1.7 g (10 mmol)) synthesized in Reference Example 1 and 0.6 g (15 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Furthermore, 7.1 g (10 mmol) of an aqueous 10% sodium thiomethoxide solution was added dropwise thereto at room temperature, and the obtained mixture was stirred for 6 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, and the precipitated crystal was collected by filtration, washed twice with 5 mL of water and dried by a hot air drier to obtain 1.6 g (yield: 72.7%) of the title compound as a pale yellow crystal. This crystal was recrystallized from water-methanol and obtained as a white crystal.

20 ¹H-NMR value (300 MHz, MeOH-d₄): σ=4.86 (br, 1H), 3.64 (s, 3H), 3.56 (s, 2H), 2.02 (s, 3H) ppm.

30 LC-MS (EI): m/z=226 (M⁺), 179 (base), melting point: 123-124°C.

35 Example 2: Synthesis of [(5-hydroxy-1-methyl-3-trifluoromethylpyrazol-4-yl)-methylthio]benzene

5-Hydroxy-1-methyl-3-trifluoromethylpyrazole (1.7 g (10 mmol)) synthesized in Reference Example 1 and 0.6 g

(15 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Furthermore, 11.0 g (10 mmol) of thiophenol was added dropwise thereto at room temperature, and the obtained mixture was stirred for 7 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, and the precipitated crystal was collected by filtration, washed twice with 5 mL of water and dried by a hot air drier to obtain 2.2 g (yield: 76.4%) of the title compound as a pale yellow crystal. This crystal was recrystallized from n-hexane-2-propanol and obtained as a white crystal.

¹H-NMR value (300 MHz, MeOH-d₄): σ =7.1-7.3 (m, 5H), 4.86 (br, 1H), 3.99 (s, 2H), 3.61 (s, 3H).

LC-MS (EI): m/z=288 (M⁺), 110 (base), melting point: 152°C.

Example 3: Synthesis of 3-[(5-hydroxy-1-methyl-3-trifluoromethylpyrazol-4-yl)-methylthio]-4,5-dihydro-5,5-dimethylisooxazole

5-Hydroxy-1-methyl-3-trifluoromethylpyrazole (1.7 g (10 mmol)) synthesized in Reference Example 1 and 1.6 g (40 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Furthermore, a water 10 mL solution containing 2.1 g (10 mmol) of [5,5-dimethyl(4,5-dihydroisooxazol-3-yl)]thiocarboxamidine hydrochloride was added dropwise thereto at room temperature, and the obtained mixture was stirred at the same temperature for 2 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, and the precipitated crystal was collected by filtration, washed twice with 5 mL of water and dried by a hot air drier to

obtain 2.5 g (yield: 80.1%) of the title compound as a pale yellow crystal. This crystal was recrystallized from n-hexane-2-propanol and obtained as a white crystal.

¹H-NMR value (300 MHz, MeOH-d₄): σ =4.88 (br, 1H),
5 4.08 (s, 2H), 3.64 (s, 3H), 2.91 (s, 2H), 1.39 (s, 6H)
ppm.

LC-MS (EI): m/z=309 (M⁺), 177 (base), melting point:
115-116°C.

10 Example 4: Synthesis of 3-[(5-hydroxy-1-methyl-3-
trifluoromethylpyrazol-4-yl)-methylthio]-4,5-dihydro-5,5-
dimethylisooxazole

5-Hydroxy-1-methyl-3-trifluoromethylpyrazole (1.7 g
(10 mmol)) synthesized in Reference Example 1 and 6.0 g
15 (30 mmol) of a methanol solution of 28% sodium methoxide
were dissolved in 10 mL of methanol. While stirring the
resulting solution at room temperature, 1.7 g of
paraformaldehyde was charged thereto, followed by
stirring at the same temperature for 1 hour.
20 Furthermore, 2.1 g (10 mmol) of [5,5-dimethyl(4,5-
dihydroisooxazol-3-yl)]thiocarboxamidine hydrochloride
was charged thereto at room temperature, and the obtained
mixture was stirred for 2 hours. After the reaction, 5.0
g (50 mmol) of 35% hydrochloric acid was added dropwise
25 thereto, 10 mL of water was further added, and the
precipitated crystal was collected by filtration, washed
twice with 5 mL of water and dried by a hot air drier to
obtain 2.6 g (yield: 84.1%) of the title compound as a
pale yellow crystal. The ¹H-NMR spectrum agreed with that
30 of Example 3.

Example 5: Synthesis of 3-[(5-hydroxy-1-methyl-3-
trifluoromethylpyrazol-4-yl)-methylthio]-4,5-dihydro-5,5-
dimethylisooxazole

35 The title compound (2.3 g (yield: 74.2%)) was
obtained as a pale yellow crystal by the same operation
as in Example 4 except for changing the base to 4.2 g (30

mmol) of potassium carbonate in the reaction of Example 4. The ¹H-NMR spectrum agreed with that of Example 3.

Example 6 (embodiment where the electron-withdrawing group is a trifluoromethyl group): Synthesis of 4-[(5-hydroxy-1-methyl-3-trifluoromethylpyrazol-4-yl)-methylsulfonyl]toluene

5-Hydroxy-1-methyl-3-trifluoromethylpyrazole (8.3 g (50 mmol)) synthesized in Reference Example 1 and 3.0 g (75 mmol) of sodium hydroxide were dissolved in 50 mL of water. While stirring the resulting solution at room temperature, 8.5 g (100 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Furthermore, 9.0 g (50 mmol) of sodium p-toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 25.0 g (250 mmol) of 35% hydrochloric acid was added dropwise thereto, 100 mL of water was further added, and the precipitated crystal was collected by filtration, washed twice with 20 mL of water and dried by a hot air drier to obtain 14.0 g (yield: 83.8%) of the title compound as a white crystal.

¹H-NMR value (300 MHz, MeOH-d₄): σ =7.62 (d; J=8.4 Hz, 2H), 7.39 (d; J=8.4 Hz, 2H), 4.85 (br, 1H), 4.32 (s, 2H), 3.63 (s, 3H), 2.44 (s, 3H) ppm.

LC-MS (EI): m/z=334 (M⁺), 179 (base), melting point: 135°C.

Example 7: Synthesis of 4-[(5-hydroxy-1-methyl-3-trifluoromethylpyrazol-4-yl)-methylsulfonyl]toluene

5-Hydroxy-1-methyl-3-trifluoromethylpyrazole (1.7 g (10 mmol)) synthesized in Reference Example 1 and 0.6 g (15 mmol) of sodium hydroxide were dissolved in 10 mL of DMF. While stirring the resulting solution at room temperature, 1.7 g of paraformaldehyde was charged thereto, followed by stirring at the same temperature for 1 hour. Furthermore, 1.8 g (10 mmol) of sodium p-

toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, 10 mL of water was further added, and the precipitated crystal was collected by filtration, washed twice with 5 mL of water and dried by a hot air drier to obtain 3.0 g (yield: 88.2%) of the title compound as a white crystal. The ¹H-NMR spectrum agreed with that of Example 6.

10

Example 8: Synthesis of 4-[(5-hydroxy-1-methyl-3-trifluoromethylpyrazol-4-yl)-methylsulfonyl]toluene

5-Hydroxy-1-methyl-3-trifluoromethylpyrazole (1.7 g (10 mmol)) synthesized in Reference Example 1 and 0.6 g (15 mmol) of sodium hydroxide were dissolved in 10 mL of toluene. While stirring the resulting solution at room temperature, 1.7 g of paraformaldehyde was charged thereto, followed by stirring at the same temperature for 24 hours. Furthermore, 1.8 g (10 mmol) of sodium p-toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 8 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, 30 mL of water was further added, and the precipitated crystal was collected by filtration, washed twice with 5 mL of water and dried by a hot air drier to obtain 3.0 g (yield: 88.2%) of the title compound as a pale yellow crystal. The ¹H-NMR spectrum agreed with that of Example 6.

30 Example 9: Synthesis of 3-[(5-hydroxy-1-phenyl-3-trifluoromethylpyrazol-4-yl)-methylthio]-4,5-dihydro-5,5-dimethylisooxazole

5-Hydroxy-1-phenyl-3-trifluoromethylpyrazole (2.3 g (10 mmol)) synthesized in Reference Example 2 and 0.6 g (15 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution

was added dropwise thereto, followed by stirring at the same temperature for 1 hour (Reaction Solution 1). Separately, 2.1 g (10 mmol) of [5,5-dimethyl(4,5-dihydroisooxazol-3-yl)]thiocarboxamidine hydrochloride was added to a water 10 mL solution containing 1.2 g (30 mmol) of sodium hydroxide, followed by stirring for 1 hour (Reaction Solution 2). Reaction Solution 2 was added dropwise to Reaction Solution 1, and the obtained mixture was stirred for 2 hours. After the reaction, 6.0 g (60 mmol) of 35% hydrochloric acid was added dropwise thereto, and the toluene layer obtained from an extraction operation performed twice with 20 mL of toluene was washed with 10 mL of water and then with 10 mL of saturated brine and dried over anhydrous sodium sulfate. Thereafter, toluene was removed by distillation under reduced pressure to obtain 3.3 g (yield: 89.2%) of the title compound as a viscous material. This viscous material was left standing at room temperature for 2 days and thereby crystallized.

¹H-NMR value (300 MHz, MeOH-d₄): σ =7.3-7.7 (m, 4H), 4.90 (br, 1H), 4.16 (s, 2H), 2.92 (s, 2H), 1.37 (s, 6H) ppm.

Melting point: 89-92°C.

Example 10: Synthesis of 4-[(5-hydroxy-1-phenyl-3-trifluoromethylpyrazol-4-yl)-methylsulfonyl]benzene

5-Hydroxy-1-phenyl-3-trifluoromethylpyrazole (2.3 g (10 mmol)) synthesized in Reference Example 2 and 0.6 g (15 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Subsequently, 1.6 g (10 mmol) of sodium benzenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 6.0 g (60 mmol) of 35% hydrochloric acid was added dropwise thereto, and the

toluene layer obtained from an extraction operation performed twice with 20 mL of toluene was washed with 10 mL of water and then with 10 mL of saturated brine and dried over anhydrous sodium sulfate. Thereafter, toluene was removed by distillation under reduced pressure to obtain 3.4 g (yield: 88.1%) of the title compound as a viscous material. This viscous material was left standing at room temperature for 2 days and thereby crystallized.

^1H -NMR value (300 MHz, MeOH- d_4): σ =7.3-7.9 (m; 10 H), 4.91 (br, 1H), 4.44 (s, 2H), 3.63 (s, 3H), 2.44 (s, 3H) ppm.

Melting point: 122-123°C.

Example 11: Synthesis of 3-[(5-hydroxy-3-trifluoromethylpyrazol-4-yl)-methylthio]-4,5-dihydro-5,5-dimethylisooxazole

5-Hydroxy-3-trifluoromethylpyrazole hydrochloride (1.9 g (10 mmol)) synthesized in Reference Example 3 and 1.2 g (30 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour (Reaction Solution 1). Separately, 2.1 g (10 mmol) of [5,5-dimethyl(4,5-dihydroisooxazol-3-yl)]thiocarboxamidine hydrochloride was added to a water 10 mL solution containing 1.2 g (30 mmol) of sodium hydroxide, followed by stirring for 1 hour (Reaction Solution 2). (Reaction Solution 2) was added dropwise to (Reaction Solution 1), and the obtained mixture was stirred for 2 hours. After the reaction, 6.0 g (60 mmol) of 35% hydrochloric acid was added dropwise thereto, and sodium hydrogencarbonate was then added carefully so that the reaction solution could have a pH of 7. The toluene layer obtained from an extraction operation performed twice with 20 mL of ethyl acetate was washed with 10 mL of water and then with 10 mL of

saturated brine and dried over anhydrous sodium sulfate. Thereafter, ethyl acetate was removed by distillation under reduced pressure to obtain 2.4 g (yield: 81.3%) of the title compound as a viscous material. This viscous material was left standing at room temperature for 2 days and thereby crystallized.

^1H -NMR value (300 MHz, MeOH- d_4): σ =5.35 (s; 1H), 4.86 (br, 1H), 4.10 (s, 2H), 2.92 (s, 2H), 1.39 (s, 6H) ppm.

LC-MS (EI): m/z =320 (M^+), 163 (base), melting point: 131-133°C.

Example 12: Synthesis of 3-[(5-hydroxy-3-trifluoromethylpyrazol-4-yl)-methylsulfonyl]toluene

5-Hydroxy-3-trifluoromethylpyrazole hydrochloride (1.9 g (10 mmol)) synthesized in Reference Example 3 and 1.2 g (30 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Subsequently, 1.8 g (10 mmol) of sodium p-toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, 20 mL of water was further added, and the precipitated crystal was collected by filtration, washed twice with 10 mL of water and dried by a hot air drier to obtain 2.67 g (yield: 83.2%) of the title compound as a pale yellow crystal.

^1H -NMR value (300 MHz, MeOH- d_4): σ =7.62 (d; J =8.1 Hz, 2H), 7.37 (d; J =8.1 Hz, 2H), 5.34 (s; 1H), 4.88 (s, 1H), 4.34 (s, 2H), 2.43 (s, 3H) ppm.

LC-MS (EI): m/z =295 (M^+), 163 (base), melting point: 130-133°C.

Example 13: Synthesis of 4-[(3-cyano-5-hydroxy-1-

phenylpyrazol-4-yl)-methylsulfonyl]toluene

3-Cyano-5-hydroxy-1-phenylpyrazole (1.8 g (10 mmol)) synthesized in Reference Example 5 and 0.6 g (15 mmol) of sodium hydroxide were dissolved in 10 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Subsequently, 1.8 g (10 mmol) of sodium p-toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, 20 mL of water was further added, and the precipitated crystal was collected by filtration, washed twice with 20 mL of water and dried by a hot air drier to obtain 3.0 g (yield: 85.7%) of the title compound as a pale yellow crystal.

$^1\text{H-NMR}$ value (300 MHz, MeOH- d_4): σ =7.4-7.7 (m; 9H), 4.86 (s, 1H), 4.40 (s, 2H), 2.46 (s, 3H) ppm.

LC-MS (EI): m/z =353 (M^+), 197 (base), melting point: 214°C.

Example 14: Synthesis of 4-[(3-carboxy-5-hydroxy-1-phenylpyrazol-4-yl)-methylsulfonyl]toluene

5-Hydroxy-1-methyl-3-ethoxycarbonylpyrazole (1.7 g (10 mmol)) synthesized in Reference Example 4 and 1.2 g (30 mmol) of sodium hydroxide were dissolved in 20 mL of water. While stirring the resulting solution at room temperature, 1.7 g (20 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Subsequently, 1.8 g (10 mmol) of sodium p-toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 5.0 g (50 mmol) of 35% hydrochloric acid was added dropwise thereto, 30 mL of water was further added, and the precipitated crystal was collected by filtration, washed twice with 20 mL of water and dried by a hot air drier to obtain 2.6 g (yield:

82.9%) of the title compound as a white crystal.

¹H-NMR value (300 MHz, MeOH-d₄): σ =7.52 (d; J=8.1 Hz, 2H), 7.33 (d; J=8.1 Hz, 2H), 4.85 (s, 2H), 4.61 (s, 2H), 3.65 (s, 3H), 2.42 (s, 3H) ppm.

5 LC-MS (EI): m/z=310 (M⁺), 155 (base), melting point: 228°C.

Example 15: Synthesis of 4-[[3-(2,4-dichlorophenyl)-5-hydroxy-1-methylpyrazol-4-yl]-methylsulfonyl]toluene

10 3-(2,4-Dichlorophenyl)-5-hydroxy-1-methylpyrazole (1.3 g (5 mmol)) synthesized in Reference Example 6 and 0.3 g (7.5 mmol) of sodium hydroxide were dissolved in 5 mL of ethanol. While stirring the resulting solution at room temperature, 1.0 g (11 mmol) of a 35% formalin
15 solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour. Subsequently, 0.9 g (5 mmol) of sodium p-toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 2.5 g (25 mmol) of 35%
20 hydrochloric acid was added dropwise thereto, 15 mL of water was further added, and the precipitated crystal was collected by filtration, washed twice with 10 mL of water and dried by a hot air drier to obtain 2.1 g (yield: 90.3%) of the title compound as a pale yellow crystal.

25 ¹H-NMR value (300 MHz, CDCl₃): σ =7.0-7.4 (m, 7H), 4.24 (s, 2H), 3.69 (s, 3H), 3.50 (br, 1H), 2.43 (s, 3H) ppm.

LC-MS (EI): m/z=410 [(M-1)⁺], 255 (base), melting point: 209°C.

30

Example 16: Synthesis of 4-[[3-(3,5-dinitrophenyl)-5-hydroxy-1-methylpyrazol-4-yl]-methylsulfonyl]toluene

35 3-(3,5-Dinitrophenyl)-5-hydroxy-1-methylpyrazole (0.65 g (2.5 mmol)) synthesized in Reference Example 4 and 0.15 g (3.8 mmol) of sodium hydroxide were dissolved in 5 mL of ethanol. While stirring the resulting

solution at room temperature, 0.5 g (5.5 mmol) of a 35% formalin solution was added dropwise thereto, followed by stirring at the same temperature for 1 hour.

Subsequently, 0.45 g (2.5 mmol) of sodium p-

5 toluenesulfinate was charged thereto at room temperature, and the obtained mixture was stirred for 2 hours. After the reaction, 2.5 g (25 mmol) of 35% hydrochloric acid was added dropwise thereto, 15 mL of water was further added, and the precipitated crystal was collected by
10 filtration, washed twice with 10 mL of water and dried by a hot air drier to obtain 0.9 g (yield: 89.3%) of the title compound as a brown crystal.

^1H -NMR value (300 MHz, CDCl_3): σ =8.88 (t; J=2.1 Hz, 1H), 8.52 (d; J=1.8 Hz, 2H), 7.55 (d; J=8.1 Hz, 2H), 7.16
15 (d; J=8.1 Hz, 2H), 4.43 (s, 2H), 3.85 (br, 1H), 3.76 (s, 3H), 2.31 (s, 3H) ppm.

LC-MS (EI): m/z=432 (M^+), 276 (base), melting point: 192-194°C.

20 INDUSTRIAL APPLICABILITY

A novel industrial process for producing a 5-hydroxy-4-thiomethylpyrazole compound is provided. According to the process of the present invention, a 5-hydroxy-4-thiomethylpyrazole compound is produced from a
25 5-hydroxypyrazole compound represented by formula (1) in a good yield through a single step by a simple and easy operation method under mild conditions without using a special reaction apparatus or an expensive catalyst or transition metal. Moreover, a harmful waste derived from
30 a catalyst or transition metal is substantially not produced and therefore, this process is environment-friendly and has high industrial utility value.

In addition, according to the embodiment of using water as the solvent, a more environment-friendly process
35 having a higher industrial utility value is provided.